


# Cell Reports

## Anti-Müllerian Hormone Signaling Regulates Epithelial Plasticity and Chemoresistance in Lung Cancer

### Graphical Abstract



### Authors

Tim N. Beck, Vladislav A. Korobeynikov, Alexander E. Kudinov, ..., David A. Proia, Ilya G. Serebriiskii, Erica A. Golemis

### Correspondence

erica.golemis@fccc.edu

### In Brief

Beck et al. identify active signaling by the TGF- $\beta$ /BMP superfamily member anti-Müllerian hormone (AMH) and its receptor AMHR2 in non-small cell lung cancer (NSCLC), demonstrating a role for AMH/AMHR2 in influencing the basal and BMP-dependent SMAD signaling that constrains epithelial-mesenchymal transition (EMT) and in regulating drug resistance.

### Highlights

- TGF- $\beta$  superfamily member AMH regulates tumor growth and drug resistance in NSCLC
- AMH and AMHR2 activity influences SMAD, AKT, and NF- $\kappa$ B signaling in NSCLC cells
- Loss of AMH/AMHR2 promotes EMT through direct modulation of TGF- $\beta$ /BMP receptors
- EMT promotes chemoresistance, but sensitizes NSCLC cells to HSP90 inhibition



Beck et al., 2016, Cell Reports 16, 657–671  
July 19, 2016 © 2016 The Author(s).  
<http://dx.doi.org/10.1016/j.celrep.2016.06.043>

CellPress